

1 **SOURCES AND CONCENTRATIONS OF CUPRIC FUNGICIDES FOR THE**  
2 **CONTROL OF CITRUS BLACK SPOT<sup>1</sup>**

[RC1] Comentário: Título no máximo 15 palavras.

3  
4  
5 **ABSTRACT** – Citrus black spot (CBS) is a severe disease for citriculture in the São Paulo  
6 State, Brazil. Part of its management is focused on chemical control using cupric fungicides  
7 and strobilurins. The objective of the present work was to evaluate the efficacy of three  
8 sources and three concentrations of cupric fungicides (copper hydroxide, copper oxychloride  
9 and cuprous oxide). Orange fruits of the Pera cultivar were bagged in the plants and the  
10 treatment with cupric fungicide was applied. The fruits were inoculated (by spray) with  
11 *Phyllosticta citricarpa* ( $1 \times 10^4$  conidia mL<sup>-1</sup>) after 0, 7, 14, 21, and 28 days, and bagged again.  
12 The evaluation of incidence and severity was did at the harvest time of fruits. A second  
13 experiment was conducted under natural infection with the same treatments, consisting of  
14 application of fungicides at the stages F1 (petal fall) and F2 (fruits with diameter of 1 cm),  
15 using mechanized sprayer with mean flow of 7.35 L solution plant<sup>-1</sup>. Subsequently, all  
16 treatments received four applications of azoxystrobin (30 g ha<sup>-1</sup>), plus mineral oil at 0.25%.  
17 Four monthly evaluations were done to determine the CBS incidence and severity. The initial  
18 applications with cupric fungicides are essential for the control of CBS; the fungicide copper  
19 hydroxide showed the best control of CBS with the lowest rate of metallic copper (43.7 g of  
20 Cu<sup>++</sup> 100 L<sup>-1</sup>) in both experiments, regardless of the conduction conditions.

[RC2] Comentário: Na primeira versão do artigo submetido, os nomes dos autores e a nota de rodapé com os endereços deverão ser omitidos.

21  
22 **Keywords:** *Citrus sinensis*. *Phyllosticta citricarpa*. Metallic copper.

[RC3] Comentário: Abstract no máximo 250 palavras.

[RC4] Comentário: A primeira letra maiúscula. Devem ter, no mínimo, três e, no máximo, cinco palavras, não constantes no Título/Title e separadas por ponto.

23  
24  
25 **FONTES E CONCENTRAÇÕES DE FUNGICIDAS CÚPRICOS NO CONTROLE DA**  
26 **MANCHA PRETA DOS CITROS**

[RC5] Comentário: Título no máximo 15 palavras.

27  
28  
29 **RESUMO** – A mancha preta dos citros é uma doença severa para a citricultura do Estado de  
30 São Paulo. Parte do seu manejo é focado no controle químico usando fungicidas cúpricos e  
31 estrobilurinas. Este trabalho teve o objetivo de avaliar a eficácia de três fontes e três  
32 concentrações de fungicidas cúpricos (hidróxido de cobre, oxiclureto de cobre e óxido  
33 cuproso). Frutos de laranja 'Pera' foram ensacados nas plantas seguido de tratamento com  
34 fungicida cúprico. Posteriormente, após zero; sete; 14; 21 e 28 dias tais frutos foram

35 inoculados com *Phyllosticta citricarpa* (1x10<sup>4</sup> conídios mL<sup>-1</sup>), por pulverização, seguido de  
36 novo ensacamento. A avaliação da incidência e severidade da doença ocorreu na colheita dos  
37 frutos. Um segundo experimento, sob infecção natural, foi constituído pelos mesmos  
38 tratamentos, sendo os fungicidas aplicados nos estádios F1 (queda de pétalas) e F2 (frutos  
39 com 1cm de diâmetro) mediante pulverização tratorizada e vazão média 7,35 L calda planta<sup>-1</sup>.  
40 Subsequentemente, e de forma semelhante para todos os tratamentos, foram realizadas quatro  
41 aplicações de azoxistrobina (30 g de i.a./ha) acrescida de óleo mineral a 0,25%. Foram  
42 realizadas quatro avaliações mensais para determinação da incidência e severidade dos  
43 sintomas. Concluiu-se que: as pulverizações iniciais com fungicidas cúpricos são  
44 fundamentais para o controle da mancha preta dos citros; independente das condições de  
45 condução, para ambos os ensaios o fungicida hidróxido de cobre propiciou a melhor resposta  
46 de controle da mancha preta dos citros com a menor dosagem de cobre metálico (43,7 g de  
47 Cu<sup>++</sup>100L<sup>-1</sup>).

[RC6] Comentário: Resumo no máximo 250 palavras.

48  
49 **Palavras-chave:** *Citrus sinensis*. *Phyllosticta citricarpa*. Cobre metálico.

[RC7] Comentário: A primeira letra maiúscula. Devem ter, no mínimo, três e, no máximo, cinco palavras, não constantes no Título/Title e separadas por ponto.

## 52 INTRODUCTION

53  
54 **Citrus** black spot (CBS) is caused by the fungus *Phyllosticta citricarpa* McAlp.  
55 (teleomorph: *Guignardia citricarpa* Kiely) (BALDASSARI; WICKERT; GOES, 2008), and  
56 is associated with citrus plants in several countries in Africa, Asia, Oceania (KOTZÉ, 2000;  
57 EPPO, 2017), South America and North America, and in Caribe (TIMMER et al., 2000;  
58 SCHUBERT et al., 2010; HIDALGO and PÉREZ, 2010). All citrus varieties of economic  
59 importance are susceptible to this fungus, with losses that can reach 40% of the production  
60 (SILVA JÚNIOR et al., 2016).

[RC8] Comentário: Parágrafos: primeira linha 1 cm.

Especial:  Por:

61 The control of CBS is done usually by using fungicides; in Brazil, it is controlled  
62 mainly with use of cupric and strobilurin fungicides (MOTTA, 2009; VINHAS, 2011; SILVA  
63 JÚNIOR et al., 2016). Cupric fungicides are commonly applied after petal fall, corresponding  
64 to the stages F1 and F2, until the fruits reach diameter of 1 cm (STOLLER, 2010;), followed  
65 by two to five applications of strobilurin fungicides (SCALLOPPI et al., 2012), covering the  
66 more susceptible period for fruits (AGUIAR et al., 2012), from petal fall until the end of the  
67 summer, when rainfall periods of more than 8 hours still occur. Ikeda (2011) reported

[RC9] Comentário: Veja as normas de citações em: <https://periodicos.ufersa.edu.br/index.php/caatinga/preparodosmanuscritos>.

Com 1(um) autor, usar Torres (2008) ou (TORRES, 2008); com 2 (dois) autores, usar Torres e Marcos Filho (2002) ou (TORRES; MARCOS FILHO, 2002); com 3 (três) autores, usar França, Del Grossi e Marques (2009) ou (FRANÇA; DEL GROSSI; MARQUES, 2009); com mais de três autores, usar Torres et al. (2002) ou (TORRES et al., 2002).

68 satisfactory responses of control of CBS to the use of five or six applications of cupric  
69 fungicides combined or alternated with strobilurins.

70 Cupric fungicides are approved, for citrus plants, only for the control of citrus scab and  
71 melanosis, except copper hydroxide (Kocide WDG Bioactive, Mitsui & Co. Brasil, S.A.)  
72 (MAPA, 2017); however, they are used isolate or in combination with strobilurins for the  
73 control of *P. citricarpa* (VINHAS, 2011). These fungicides are applied with four-week  
74 intervals, beginning after  $\frac{3}{4}$  of petals felled, and subsequently with 28-day intervals,  
75 combined with application of strobilurins at 35-day to 42-day intervals (MOTTA, 2009). The  
76 recommended rates of cupric fungicides are 0.75 to 1.25 g L<sup>-1</sup> for copper hydroxide, 2.5 g L<sup>-1</sup>  
77 for copper oxychloride, and 1.0 g L<sup>-1</sup> for cuprous oxide (MAPA, 2017). However, the rates  
78 used at the stages F1 and F2, and in subsequent applications in combination with strobilurins  
79 for the control of CBS under field conditions vary, and are usually based on the fungicide  
80 metallic copper contents. These recommendations are from studies with copper oxychloride;  
81 no studies comparing sources and concentrations of cupric fungicides are found.

82 In this context, the objective of the present work was to evaluate the efficacy of three  
83 sources and three concentrations of cupric fungicides under artificial conditions of inoculation  
84 with *P. citricarpa*, and under natural conditions, to determine the equivalence between the  
85 different compounds in relation to metallic copper concentration in the solution.

86  
87

## 88 MATERIAL AND METHODS

89

### 90 Experiment I – Protector effect for different sources and rates of cupric fungicides 91 against *Phyllosticta citricarpa* infection in orange fruits of the Pera cultivar

92

93 The experiment was conducted in 2012 in an orchard with orange plants of the Pera  
94 cultivar (plant spacing of 5.5 × 2 m), at a private property in Olímpia, state of São Paulo (SP),  
95 Brazil (20°41'57.23"S and 48°59'35.33"W).

96 The isolated pathogen was from Conchal, SP, Brazil. It was multiplied in Petri dishes  
97 containing BDA medium and maintained in BOD at temperature of 25°C.

98 The fungicides evaluated were copper hydroxide (Kocide WDG Bioactive<sup>®</sup> 462 g of  
99 copper hydroxide, Mitsui & Co. Brasil, S.A.), copper oxychloride (Recop<sup>®</sup> 840 g of copper  
100 oxychloride, Atanor do Brasil), and cuprous oxide (Redshield 750<sup>®</sup> 750 g of cuprous oxide,

[RC10] Comentário: Não pode separar este tópico.

101 Agrovant Comércio de Produtos Agrícolas Ltda). The treatments and rates used are shown in  
 102 Table 1.

103 Approximately 5,000 fruits at stage F2 (STOLLER, 2010) of 100 plants were bagged on  
 104 02/23/2015 to avoid infections by *P. citricarpa* naturally presents in the area. When the plants  
 105 were at stage F4, they were treated with cupric fungicide on 05/13/2016. Subsequently, the  
 106 fruits were inoculated with *P. citricarpa* ( $1 \times 10^4$  conidia mL<sup>-1</sup>) in intervals of 0, 7, 14, 21, and  
 107 28 days. The inoculum was prepared and the inoculation was done according to the  
 108 methodology described by Almeida et al. (2008). After inoculation, the fruits were bagged  
 109 again, using double layer bags (crystal paper), according to the methodology adopted by  
 110 Motta (2009). The fruits were kept in the bag until the harvest (10/31/2016).

111  
 112 **Table 1.** Description of the cupric fungicide treatments for the control of citrus black spot in  
 113 orange plants of the Pera cultivar, and their active ingredient concentrations, formulated  
 114 product rates, and metallic copper equivalent rates.

**[RC11] Comentário:** Não é aceito tabela e/ou figura com ORIENTAÇÃO na forma “paisagem”, ou que apresentem mais de 17 cm de largura.

Treatments	Concentration g kg <sup>-1</sup>	Rate per 100 liters of solution	
		Formulated Product	Metallic Copper
Copper hydroxide (538 g kg <sup>-1</sup> )	538	125.0	43.7
Copper hydroxide (538 g kg <sup>-1</sup> )	538	100.0	35.0
Copper hydroxide (538 g kg <sup>-1</sup> )	538	75.0	26.2
Copper oxychloride (840 g kg <sup>-1</sup> )	840	200.0	100.8
Copper oxychloride (840 g kg <sup>-1</sup> )	840	180.0	90.7
Copper oxychloride (840 g kg <sup>-1</sup> )	840	135.0	68.0
Cuprous oxide (860 g kg <sup>-1</sup> )	860	120.0	90.0
Cuprous oxide (860 g kg <sup>-1</sup> )	860	75.0	56.3
Cuprous oxide (860 g kg <sup>-1</sup> )	860	56.3	42.2
Control			

115 Copper hydroxide = Kocide WDG Bioactive<sup>®</sup>, Mitsui & Co. Brasil, S.A.; Copper oxychloride  
 116 = Recop<sup>®</sup>, Atanor do Brasil; and Cuprous oxide = Redshield 750<sup>®</sup>, Agrovant Comércio de  
 117 Produtos Agrícolas Ltda.

118

119 The fungicides were applied according to an adaption of the method of Motta (2009);  
120 fruits with diameter of 40 to 50 mm were individually sprayed with fungicide, plus mineral oil  
121 at 0.20% (v v<sup>-1</sup>).

122 A completely randomized experimental design was used, with a 3×3×5 factorial  
123 arrangement consisted of three fungicides, three rates, and 5 inoculation times, with four  
124 replications. The plots consisted of 20 fruits. The control treatment consisted of fruits without  
125 application of cupric fungicide and with artificial inoculation with *P. citricarpa*.

126 The evaluations consisted of determination of the incidence (%) of symptomatic fruits  
127 for citrus black spot (CBS), and the disease severity. The CBS severity was estimated using a  
128 scale of grades, according to visual symptoms in percentage of lesion area—0 = fruits without  
129 visual symptoms, 1= up to 0.8%, 2 = 0.8% to 1.6%, 3 = 1.6% to 3.1%, 4 = 3.1% to 6.2%, 5 =  
130 6.2% to 12.5%, 6 = more than 6.2% (SPÓSITO et al., 2004).

131 The grades attributed to fruits were used to determine the disease index (DI), according  
132 to Wheeler (1969):

$$DI = \frac{1}{N} \sum_{i=0}^m i \cdot n_i$$

133 where  $N$  is the number of total fruits evaluated,  $i$  is the grade of the fruit,  $n_i$  is the number of  
134 fruits with grade  $I$ , and  $m$  is the maximum grade.

135 The grades attributed, disease index, and the other parameters were subjected to  
136 analysis of variance, and the means were compared by the Scott-Knott test at 5% probability  
137 of error.

138

## 139 **Experiment II - Evaluation of rates of cupric fungicides for the control of citrus black** 140 **spot**

141

142 The experiment was conducted in 2007, under natural conditions of infection by *P.*  
143 *citricarpa*, in an orchard with orange plants of the Pera cultivar (plant spacing of 7 × 3.5 m),  
144 at a private property of commercial production in Olímpia, SP, Brazil (20°47'59.17"S,  
145 49°2'34.94"W).

146 The treatments evaluated in this experiment, representing the cupric fungicide  
147 variations, were the same used in Experiment I (Table 1).

148 The fungicides were applied using a mechanized sprayer (FM Copling) with 54 nozzles  
149 MagnoJet-DDC4, diffusers 25 with 758.42 kPa at 540 RPM, power takeoff with 1900 RPM,

150 and tractor speed of 3.4 km h<sup>-1</sup>. The mean flowrate was 7.35 L plant<sup>-1</sup>, equivalent to 134 mL  
151 m<sup>-3</sup> on the plants' canopies. The cupric fungicides were sprayed when the plants were at the  
152 stages F1 and F2 (STOLLER, 2010), on 12/02/2015 and 12/23/2015. The spraying of  
153 fungicides were resumed on 01/13/2016 with foliar applications of 500 g of azoxystrobin  
154 (Vantigo<sup>®</sup>, Syngenta Proteção de Cultivos Ltda, São Paulo) at concentration of 16 g per 100 L  
155 of water plus mineral oil (Agefix<sup>®</sup>, Packblend Indústria e Comércio de Lubrificantes Ltda) at  
156 0.25% (v v<sup>-1</sup>) until 05/19/2016, totaling four applications with 42-day intervals.

157 A randomized block experimental design was used, consisting of 10 treatments and four  
158 replications. The plots consisted of three rows of 11 plants, totaling 33 plants or 808.5 m<sup>2</sup>.

159 The evaluations were carried out on 08/10/2016, 09/08/2016, 10/06/2016, and  
160 11/14/2016 to determine the CBS incidence and severity in 100 random fruits collected from  
161 the five central plants of each plot. The CBS severity was estimated using a scale of grades  
162 (SPÓSITO et al., 2004), and the data were used to determine the disease index (DI),  
163 according to the same equation used in Experiment I.

164 The DI data were used to determine the area under the disease progress curve (AUDPC)  
165 by the trapezoidal method  $(DI1 + DI2)/2 * (Day2 - Day1)$ . The fruits were harvested on  
166 11/30/2016, when the fruit yield per plant (kg) was determined. The data were subjected to  
167 analysis of variance and the means were compared by the Scott-Knott test at 5% probability  
168 of error. The data were also subjected to regression analysis to determine the rates of each  
169 fungicide based on linear and quadratic responses of the AUDPC.

170  
171

## 172 **RESULTS AND DISCUSSION**

173

### 174 **Experiment I**

175

176 All treatments with the highest rates of cupric fungicides presented on average lower  
177 incidence and severity of citrus black spot (CBS) (Table 2). The results showed direct and  
178 significant responses between fungicide rates and CBS incidence and severity, except for the  
179 treatment with copper oxychloride at 90.7 g 100 L<sup>-1</sup>. However, high rates on equivalent  
180 metallic copper do not necessarily mean high concentrations of fungicide (SILVA JÚNIOR et  
181 al., 2016).

182 The number of days after the application of fungicides had a significant and negative  
183 effect, with lower CBS incidence and severity when the inoculation with *Phyllosticta*

[RC12] Comentário: Não pode separar este tópico.

184 *citricarpa* and the treatment was carried out on the same day. The inoculations between 7 and  
 185 28 days after the fungicide application resulted in similar protections. This result is related to  
 186 those reported by Motta (2009), who found incidence of 30% and 80% when the inoculation  
 187 was at 1 and 28 days after application of copper oxychloride, respectively. Thus, the  
 188 methodology used is efficient to evaluate and replicate results, and can be used to evaluate  
 189 new formulations or fungicides.

190 The interaction between the fungicide rates, fungicide sources, and days after treatment  
 191 of fruits was not significant, denoting that the protection tends to decrease over time at the  
 192 same proportion for all the treatments, even when using different fungicide rates and sources.  
 193 These results showed that the best interval between applications of cupric fungicides depends  
 194 on the control level desired, in terms of CBS incidence and severity. Thus, it can be weekly  
 195 for production of fruits with lower incidence, and up to 28 days for production of fruits with  
 196 incidence and severity within limits that avoid the fall of fruits. Motta (2009) found similar  
 197 results, with increases in CBS incidence and severity as a function of increases in the intervals  
 198 between the application of copper oxychloride and artificial inoculation with *P. citricarpa*;  
 199 they reported satisfactory protection up to 21 days after the application.

200

201 **Table 2.** Incidence and disease index of citrus black spot (*Phyllosticta citricarpa*) in orange  
 202 fruits of the Pera cultivar as a function of cupric fungicides rates and sources and days after  
 203 the application, in the 2015/2016 crop season.

Sources of variation		Incidence (%)	Disease index
Treatments	Rate (g of metallic copper 100 L <sup>-1</sup> )		
1. Copper hydroxide (538 g kg <sup>-1</sup> )	43.7	16.9 A	0.22 a
2. Copper hydroxide (538 g kg <sup>-1</sup> )	35.0	24.5 b	0.31 b
3. Copper hydroxide (538 g kg <sup>-1</sup> )	26.2	26.5 b	0.34 b
4. Copper oxychloride (840 g kg <sup>-1</sup> )	100.8	14.7 a	0.19 a
5. Copper oxychloride (840 g kg <sup>-1</sup> )	90.7	13.0 a	0.17 a
6. Copper oxychloride (840 g kg <sup>-1</sup> )	68.0	32.5 c	0.41 c
7. Cuprous oxide (860 g kg <sup>-1</sup> )	90.0	17.9 a	0.24 a
8. Cuprous oxide (860 g kg <sup>-1</sup> )	56.3	20.8 b	0.27 b

9. Cuprous oxide (860 g kg <sup>-1</sup> )	42.2	38.9 c	0.49 c
10. Control		54.0 d	0.68 d
F Test		13.612 **	18.058 **
Days after application			
0		18.7 a	0.25 a
7		25.7 b	0.33 b
14		26.5 b	0.34 b
21		31.2 b	0.40 b
28		27.8 b	0.35 b
F Test		5.047 **	4.996 **
Interaction (Treatment × Days after application)			
F Test		1.052 ns	1.103 ns
CV (%)		32.88	10.87

204 The data of incidence and severity of citrus black spot were transformed according to the  
205 equations  $x' = \arcsen\sqrt{x/100}$ , and  $x' = \sqrt{x + 0.5}$ , respectively. Means followed by the  
206 same letter in the columns do not differ by the Scott-Knott test at 5% probability of error. \*\*  
207 = significant by the analysis of variance at 1% probability of error; ns = not significant by the  
208 analysis of variance at 5% probability of error.

209

## 210 Experiment II

211

212 According to the first evaluations, in August, the highest cupric fungicide rates were  
213 significantly more efficient to reduce the CBS incidence (Table 3). However, they did not  
214 necessarily represent the same metallic copper rate (Table 1), contradicting the  
215 recommendation by equivalence (SILVA JÚNIOR et al., 2016b). These data showed that the  
216 rate should be specific for each cupric fungicide source, and cannot be generalized or  
217 equalized by the equivalent metallic copper. These differences were less pronounced over the  
218 evaluations, although a high effectiveness had been maintained. The CBS incidence in the  
219 control treatment varied from 44% to 94% from the first (10/08) to the last evaluation (14/11),  
220 while in the most efficient chemical treatment it varied from 28% to 77%. These are similar  
221 results to those found by Scaloppi et al. (2012), who found that better responses are  
222 dependent on the simultaneous use of different alternatives, including fungicide with different



223 properties from protectors and cultural practices that reduce the inoculum, thus, reducing the  
 224 CBS incidence and severity.

225

226 **Table 3.** Incidence of citrus black spot (*Phyllosticta citricarpa*) in orange fruits of the Pera  
 227 cultivar treated with different cupric fungicides rates and sources in the 2015/2016 crop  
 228 season.

Treatments	Rate g 100 L <sup>-1</sup>	Incidence			
		Evaluation 1	Evaluation 2	Evaluation 3	Evaluation 4
		08/10/2016	09/08/2016	10/06/2016	11/14/2016
Copper hydroxide (538 g kg <sup>-1</sup> )	125.0	22.0 a	33.1 a	46.0 a	67.3 a
Copper hydroxide (538 g kg <sup>-1</sup> )	100.0	32.8 b	42.0 a	52.3 a	81.5 b
Copper hydroxide (538 g kg <sup>-1</sup> )	75.0	42.3 b	43.8 a	46.3 a	77.0 a
Copper oxychloride (840 g kg <sup>-1</sup> )	200.0	23.0 a	36.5 a	51.0 a	73.0 a
Copper oxychloride (840 g kg <sup>-1</sup> )	180.0	22.0 a	36.7 a	52.8 a	73.0 a
Copper oxychloride (840 g kg <sup>-1</sup> )	135.0	40.3 b	48.0 b	56.5 a	88.8 b
Cuprous oxide (860 g kg <sup>-1</sup> )	120.0	33.3 b	36.5 a	40.8 a	72.5 a
Cuprous oxide (860 g kg <sup>-1</sup> )	75.0	38.8 b	41.3 a	44.5 a	74.5 a
Cuprous oxide (860 g kg <sup>-1</sup> )	56.3	46.6 b	49.6 b	53.3 a	87.0 b
Control		44.4 b	58.4 c	72.8 b	94.3 b
F Test		6.563 **	8.11 **	3.964 **	2.92 *
CV (%)		8.40	6.40	12.31	39.10

229 The data of incidence of citrus black spot were transformed according to the equation  
 230  $x' = \arcsen\sqrt{x/100}$ . Means followed by the same letter in the columns do not differ by the  
 231 Scott-Knott test at 5% probability of error. \*\* = significant by the analysis of variance at 1%  
 232 probability of error; \* = significant by the analysis of variance at 5% probability of error.

233

234 The treatments had no effect on the CBS severity (Table 4); the CBS severity increased  
 235 even in plots treated with fungicide. No treatment completely controlled the CBS. The  
 236 fungicides and rates presented no differences for the control of CBS; they were different only  
 237 from the control treatment.

238

Treatments	Rate g 100 L <sup>-1</sup>	Severity			
		Evaluation	Evaluation	Evaluation	Evaluation
		1 08/10/2016	2 09/08/2016	3 10/06/2016	4 11/14/2016
Copper hydroxide (538 g kg <sup>-1</sup> )	125.0	0.3 a	0.4 a	0.6 a	0.9 a
Copper hydroxide (538 g kg <sup>-1</sup> )	100.0	0.4 b	0.6 b	0.8 a	1.1 a
Copper hydroxide (538 g kg <sup>-1</sup> )	75.0	0.6 d	0.6 b	0.7 a	1.2 a
Copper oxychloride (840 g kg <sup>-1</sup> )	200.0	0.2 a	0.4 a	0.6 a	0.9 a
Copper oxychloride (840 g kg <sup>-1</sup> )	180.0	0.3 a	0.5 a	0.7 a	1.0 a
Copper oxychloride (840 g kg <sup>-1</sup> )	135.0	0.4 b	0.6 b	0.8 a	1.1 a
Cuprous oxide (860 g kg <sup>-1</sup> )	120.0	0.4 b	0.4 a	0.5 a	0.9 a
Cuprous oxide (860 g kg <sup>-1</sup> )	75.0	0.5 c	0.5 b	0.6 a	1.0 a

Cuprous oxide (860 g kg <sup>-1</sup> )	56.3	0.6 d	0.6 b	0.7 a	1.1 a
Control		0.7 d	0.9 c	1.1 b	1.9 b
F Test		16.21 **	11.11 **	4.08 **	8.15 **
CV (%)		4.01	3.98	6.55	6.12

239 **Table 4.** Severity of citrus black spot (*Phyllosticta citricarpa*) in orange fruits of the Pera  
240 cultivar treated with different cupric fungicides rates and sources in the 2015/2016 crop  
241 season.

242 The data of severity of citrus black spot were transformed according to the equation  $x' =$   
243  $\sqrt{x/0.5}$ . Means followed by the same letter in the columns do not differ by the Scott-Knott  
244 test at 5% probability of error. \*\* = significant by the analysis of variance at 1% probability  
245 of error.

246  
247

248 The initial cupric fungicide applications by spraying were essential to control CBS,  
249 since the control treatment without application of cupric fungicides presented higher CBS  
250 incidence and severity (Table 5).

251 According to the area under the disease progress curve (AUDPC) for CBS incidence,  
252 only the lowest rates of copper oxychloride (135 g 100 L<sup>-1</sup>) and cuprous oxide (75 g 100 L<sup>-1</sup>)  
253 presented significant lower efficiency, increasing the CBS incidence (Table 5). The AUDPC  
254 for CBS severity showed that all the treatments were significantly different, and that they  
255 were different from the control treatment. Regarding the copper hydroxide, only the highest  
256 rate (125 g 100 L<sup>-1</sup>) was equivalent to the best treatments; the intermediate and highest rates  
257 of copper oxychloride and cuprous oxide presented the best results of control.

258

259 **Table 5.** Area under the disease progress curve (AUDPC) for incidence and severity of citrus  
260 black spot (*Phyllosticta citricarpa*) and yield of orange plants of the Pera cultivar as a  
261 function of cupric fungicides rates and sources, in the 2015/2016 crop season.

Treatments	Rate g 100 L <sup>-1</sup>	AUDPC		Orange yield
		Incidence (%)	Disease index	Kg plant <sup>-1</sup> 11/30/2016

Copper hydroxide (538 g kg <sup>-1</sup> )	125.0	4114.6 a	52.3 a	70.9 a
Copper hydroxide (538 g kg <sup>-1</sup> )	100.0	5011.8 a	68.6 b	68.6 a
Copper hydroxide (538 g kg <sup>-1</sup> )	75.0	4911.7 a	74.1 b	49.9 a
Copper oxychloride (840 g kg <sup>-1</sup> )	200.0	4504.4 a	53.7 a	75.4 a
Copper oxychloride (840 g kg <sup>-1</sup> )	180.0	4555.4 a	60.1 a	77.0 a
Copper oxychloride (840 g kg <sup>-1</sup> )	135.0	5574.4 b	71.7 b	62.3 a
Cuprous oxide (860 g kg <sup>-1</sup> )	120.0	4301.6 a	51.5 a	68.7 a
Cuprous oxide (860 g kg <sup>-1</sup> )	75.0	4682.4 a	60.8 a	61.7 a
Cuprous oxide (860 g kg <sup>-1</sup> )	56.3	5571.1 b	69.2 b	69.4 a
Control		6584.1 c	108.8 c	58.2 a
F Test		5.70 **	9.02 **	0.40 ns
CV (%)		6.17	7.88	39.46

262 The data of incidence and severity of citrus black spot were transformed according to the  
 263 equations  $x' = \arcsen\sqrt{x/100}$ , and  $x' = \sqrt{x + 0.5}$ , respectively. Means followed by the  
 264 same letter in the columns do not differ by the Scott-Knott test at 5% probability of error. \*\*  
 265 = significant by the analysis of variance at 1% probability of error; ns = not significant by the  
 266 analysis of variance at 5% probability of error.

267

268 As in Experiment I, the results of Experiment II showed no equivalence in metallic  
 269 copper rate in the treatments with the best control of CBS. The treatment 1 (copper hydroxide  
 270 at 125 g 100 L<sup>-1</sup>) had the lower metallic copper concentration in the solution (43.7 g 100 L<sup>-1</sup>).  
 271 When using 42.2 g 100 L<sup>-1</sup> with cuprous oxide, the results were significantly lower. These  
 272 were similar results to those found by Feichtenberger et al. (2001), who found equivalence of  
 273 control between cupric fungicides using a high rate of metallic copper (90 g of Cu<sup>++</sup> 100 L<sup>-1</sup>).

274 The orange yield presented no significant differences (Table 5). The sequence of the  
 275 experiment with applications of azoxystrobin maintained the CBS severity at lower levels in  
 276 the control treatment, presenting no yield losses, but losses in the fruit aesthetical quality,  
 277 which were not suited for marketing as fresh fruits.

278 Significant linear responses of the CBS were found for all fungicide treatments,  
 279 indicating that the better control levels of CBS are achieved with the use of increasing rates of  
 280 Cu<sup>++</sup> (Table 6 and Figure 1). However, this result makes unfeasible the determination of the



281 threshold for control of CBS and maximum fungicide rate, within the range of rates  
 282 established for this experiment.


283

284 **Table 6.** Regression by analysis of variance of the area under the disease progress curve  
 285 (AUDPC) for citrus black spot (*Phyllosticta citricarpa*) in orange plants of the Pera cultivar,  
 286 for each cupric fungicide as a function of the rate used, in the 2015/2016 crop season.

Cupric fungicide	Equation model	Equation	CV (%)	F Test	R <sup>2</sup>
Copper hydroxide	Linear	$y = -1.2523x + 108.79$	20.69	27.178**	98.80%
Copper oxychloride	Linear	$y = -0.5431x + 108.82$	15.38	56.837**	99.98%
Cuprous oxide	Linear	$y = -0.6483x + 103.14$	14.84	60.386**	91.85%

287





288 A = Copper hydroxide; B = Copper oxychloride; C = cuprous oxide

289 **Figure 1.** Graph of the linear regression model for the area under the disease progress curve  
 290 (AUDPC) for citrus black spot (*Phyllosticta citricarpa*) in orange of the Pera cultivar, for  
 291 each cupric fungicide as a function of the metal rate copper used, in the 2015/2016 crop  
 292 season.

293

294

## 295 CONCLUSION

296

297 The fungicides copper hydroxide, copper oxychloride, and cuprous oxide at rates of  
 298 43.7, 90.7, and 56.3 g of Cu<sup>++</sup> 100 L<sup>-1</sup>, respectively, are efficient and present similar results  
 299 for the control of citrus black spot when applied with 21-day intervals and before applications  
 300 of strobilurin fungicides.

301 The fungicides copper hydroxide, copper oxychloride, and cuprous oxide at rates of  
 302 43.7, 90.7 and 90 g of Cu<sup>++</sup> 100 L<sup>-1</sup>, respectively, can control protectively citrus fruits from  
 303 infections with *Phyllosticta citricarpa* up to 28 days.

304

305

## 306 ACKNOWLEDGEMENTS

307

308

## 309 REFERENCES

310

311 AGUIAR, R. L. et al. Período de incubação de *Guignardia citricarpa* em diferentes estádios  
 312 fenológicos de frutos de laranja 'Valência'. **Tropical Plant Pathology**, 37: 155-158, 2012.

[RC13] Comentário: As figuras devem apresentar 8,5 cm de largura, não ultrapassando 17 cm.

[RC14] Comentário: Não é aceito tabela e/ou figura com ORIENTAÇÃO na forma "paisagem", ou que apresentem mais de 17 cm de largura.

[RC15] Comentário: A resolução deve ter qualidade máxima com pelo menos 300 dpi. Não serão aceitas imagens desfocadas e com baixa qualidade.

[RC16] Comentário: Opcional.

[RC17] Comentário: Devem ser digitadas em espaço 1,5 cm e separadas entre si pelo mesmo espaço (1,5 cm).

[RC18] Comentário: O título do periódico não deve ser abreviado e recomenda-se um total de 20 a 30 referências.

[RC19] Comentário: Veja as normas de referências em <https://periodicos.ufersa.edu.br/index.php/caatinga/preparadosmanuscritos>.

313  
314 ALMEIDA, T. F.; REIS, R. F.; GOES, A. Method of inoculation of *Guignardia citricarpa*  
315 (*Phyllosticta citricarpa*) on 'Pêra Rio' sweet orange fruit. In: IX INTERNATIONAL  
316 CONGRESS OF PLANT PATHOLOGY, 2008, Torino. **Proceedings...** 2008. International  
317 Society for Plant Pathology, 2008. 1: 158-159.  
318  
319 BALDASSARI, R. B.; WICKERT, E.; GOES, A. Pathogenicity, colony morphology and  
320 diversity of isolates of *Guignardia citricarpa* and *G. mangiferae* isolated from *Citrus* spp.  
321 **Europe Journal Plant Pathology**, 120: 103-110, 2008.  
322  
323 BRASIL. Ministério da Agricultura e do Abastecimento. **AGROFIT - Sistema de**  
324 **Agrotóxicos Fitossanitários.** Disponível em:  
325 <[http://agrofit.agricultura.gov.br/agrofit\\_cons/principal\\_agrofit\\_cons](http://agrofit.agricultura.gov.br/agrofit_cons/principal_agrofit_cons)>. Acesso em: 24 mai.  
326 2017.  
327  
328 European and Mediterranean Plant Protection Organization - EPPO. **PQR-EPPO data base**  
329 **on quarantine pests.** 2017. Disponível em:  
330 <<http://www.eppo.int/DATABASES/pqr/pqr.htm>>. Acesso em: 30 ago. 2017.  
331  
332 FEICHTENBERGER, E.; et al. Competição de fungicidas à base de cobre no controle da  
333 mancha preta (*Guignardia citricarpa*) em laranjeiras 'Folha Murcha'. **Fitopatologia**  
334 **Brasileira**, 26: 444-449, 2001.  
335  
336 HIDALGO G.; PÉREZ V. L. Diferenciación morfológica, cultural y biológica de *Guignardia*  
337 *citricarpa* y *Guignardia mangiferae* en frutos cítricos de Cuba. **Fitosanidad**, 14: 141-152,  
338 2010.  
339  
340 IKEDA, M. **Efeito de fungicidas do grupo químico das estrobilurinas no controle da**  
341 **mancha preta dos citros, na produção e na qualidade tecnológica dos frutos.** 2011. 35 f.  
342 Dissertação (Mestrado em Agronomia: Área de Concentração em Produção Vegetal) -  
343 Faculdade de Ciências Agrárias e Veterinárias, Unesp, Jaboticabal, 2011.  
344

345 KOTZÉ, J. M. Citrus black spot. In: TIMMER, L.W.; GARNSEY, S. M.; GRAHAM, J. H.  
346 (Eds.). **Compendium of Citrus Diseases**. St. Paul, MN: The American Phytopathological  
347 Society, 2000. v 2, cap.6, p.595-634.  
348

349 MOTTA, R. R. **Determinação do período residual de fungicidas protetor e sistêmico para**  
350 **o controle de *Guignardia citricarpa* em frutos cítricos**. 2009. 84 f. Dissertação (Mestrado  
351 em Agronomia: Área de Concentração em Produção Vegetal) – Faculdade de Ciências  
352 Agrárias e Veterinárias, Unesp, Jaboticabal, 2009.  
353

354 SCALOPPI, E. M. T. et al. Efeito do manejo cultural e químico na incidência e severidade da  
355 mancha-preta dos citros. **Revista Brasileira de Fruticultura**, 34: 102-108, 2012.  
356

357 SCHUBERT, T. et al. **Citrus Black Spot (*Guignardia citricarpa*) Discovered in Florida**.  
358 Disponível em: <[www.doacs.state.fl.us/pi/pest\\_alerts/pdf/citrus-black-spotpest-alert.pdf](http://www.doacs.state.fl.us/pi/pest_alerts/pdf/citrus-black-spotpest-alert.pdf)>.  
359 Acesso em: 22 mai. 2017.  
360

361 SILVA JUNIOR, G. J. et al. **Pinta preta: a doença e seu manejo**. Araraquara, SP:  
362 Fundecitrus, 2016. 208 p.  
363

364 SPÓSITO, M. B. et al. Elaboração e validação de escala diagramática para avaliação da  
365 severidade da mancha preta em frutos cítricos. **Fitopatologia Brasileira**, 29: 81-85, 2004.  
366

367 STOLLER DO BRASIL LTDA. **Guia de fases de desenvolvimento: Citros**. 1. ed.  
368 Cosmópolis, SP, 2010. 4 p.  
369

370 TIMMER, L. W.; GARNSEY, S. M.; GRAHAM, J. H. **Compendium of Citrus Diseases**. 2.  
371 ed. Lake Alfred, FL: IFLA, 2000, 92 p.  
372

373 VINHAS, T. **Controle químico da *Guignardia citricarpa*, agente causal da mancha preta**  
374 **dos citros em frutos de laranja ‘Valência’**. 2011. 41 f. Dissertação (Mestrado em  
375 Agronomia: Área de Concentração em Master Citrus) - Fundo de Defesa da Citricultura,  
376 Araraquara, 2011.  
377



378 WHEELER, B. E. J. **An introduction to plant disease**. London: John Wiley & Sons, 1969.  
379 374 p.